

ENERGY FROM THE NONPERTURBATIVE QCD VACUUM

V. Gogokhia*

HAS, CRIP, RMKI, Depart. Theor. Phys., Budapest 114, P.O.B. 49, H-1525, Hungary

(Dated: February 2, 2008)

Using the effective potential approach for composite operators, we have formulated a general method of calculation of the truly nonperturbative Yang-Mills vacuum energy density (the Bag constant apart from the sign, by definition). It is the main dynamical characteristic of the QCD ground state. We define it as an integration of the truly nonperturbative effective charge over the nonperturbative region (soft momentum region). It is free of all types of the perturbative contributions, by construction. For the considered truly nonperturbative effective charge it is finite, negative and it has no imaginary part (stable vacuum), as well as it is a manifestly gauge-invariant, i.e., not explicitly depending on the gauge-fixing parameter. A nontrivial minimization procedure makes it possible to determine the Bag constant as a function either of the mass gap, which is responsible for the large-scale structure of the true QCD vacuum, or of the effective scale, which separates the nonperturbative region from the perturbative one. We have also argued that the Bag constant is a quant of the energy density which can be released from the QCD vacuum, which, in its turn, is considering as an infinite and permanent reservoir of energy.

PACS numbers: 11.15.Tk, 12.38.Lg

I. INTRODUCTION

The Lamb shift and the Casimir effect are probably the two most famous experimental evidences of zero-point energy fluctuations in the vacuum of Quantum Electrodynamics (QED) [1, 2]. The both effects are rather weak, since the QED vacuum is mainly perturbative (PT) by origin, character and magnitude (the corresponding fine structure constant is weak). However, even in this case there have been already made attempts to exploit the Casimir effect in order to release energy from the vacuum (see, for example Refs. [3, 4] and references in the above-mentioned reviews [1, 2]). Let us also note that in Ref. [5] by investigating the thermodynamical properties of the quantum vacuum it has been concluded that no energy can be extracted cyclically from the vacuum (see, however Ref. [2] and references therein).

Contrary to the QED vacuum, the vacuum of Quantum Chromodynamics (QCD) is a very complicated confining medium and its dynamical and topological complexity [6, 7] means that its structure can be organized at various levels: classical [6, 7, 8] and quantum [9]. It is mainly nonperturbative (NP) by origin, character and magnitude, since the corresponding fine structure constant is strong. So the idea to exploit the NP QCD vacuum in order to extract energy from it seems to be more attractive. However, before to discuss the ways how to extract, it is necessary to discuss which minimum/maxmimum amount of energy at all can be released in a single cycle. The vacuum of QCD due to its above-mentioned dynamical and topological complexity contains many different components and ingredients. They contribute to the truly NP vacuum energy density (VED), one of the main characteristics of the QCD ground state. It is well known that the VED in general badly diverges in quantum field theory, in particularly QCD. Thus the main problem is how to correctly define the truly NP VED which should be finite, negative and it should have no imaginary part (stable vacuum). Being one of the main physical characteristics of the QCD vacuum, the truly NP VED should not depend explicitly on the gauge-fixing parameter as well.

Let us also remind that in order to calculate physical observables from first principles in QCD it is not enough to know its Lagrangian. It is also necessary and important to know the true structure of its ground state. It is just the response of the true QCD vacuum which substantially modifies all the QCD Green's functions from their free counterparts. Just these full ("dressed") Green's functions are needed for the above-mentioned calculations.

The main purpose of this paper is twofold. First of all, to define correctly the truly NP VED, that is free of all types of the PT contributions ("PT contaminations"), as well as to make it a manifestly gauge-invariant quantity (not explicitly depending on the gauge-fixing parameter). Secondly, to discuss in general terms which amount of energy can be extracted from the QCD ground state, which in its turn is considering as an infinite and permanent reservoir of energy.

*gogohia@rmki.kfki.hu

II. THE VED

The quantum part of the VED is determined by the effective potential approach for composite operators [10, 11]. In the absence of external sources the effective potential is nothing but the VED. It is given in the form of the loops expansion, where the number of the vacuum loops (consisting in general of the confining quarks and gluons properly regularized with the help of ghosts) is equal to the power of the Plank constant, $\hbar$.

Here we are going to formulate a general method of numerical calculation of the quantum part of the truly NP Yang-Mills (YM) VED in the covariant gauge QCD. The gluon part of the VED which to-leading order (log-loop level $\sim \hbar$) is given by the effective potential for composite operators as follows [10]:

$$V(D) = \frac{i}{2} \int \frac{d^4 q}{(2\pi)^4} \text{Tr} \{ \ln(D_0^{-1} D) - (D_0^{-1} D) + 1 \}, \quad (2.1)$$

where $D(q)$ is the full gluon propagator and $D_0(q)$ is its free counterpart (see below). The traces over space-time and color group indices are understood. Evidently, the effective potential is normalized to $V(D_0) = 0$ as usual. Next-to-leading and higher contributions (two and more vacuum loops) are suppressed by one order of magnitude in powers of $\hbar$ at least, and thus are not important from the numerical point of view.

The two-point Green's function, describing the full gluon propagator, is

$$D_{\mu\nu}(q) = - \{ T_{\mu\nu}(q) d(-q^2, \xi) + \xi L_{\mu\nu}(q) \} \frac{1}{q^2}, \quad (2.2)$$

where ξ is the gauge-fixing parameter and $T_{\mu\nu}(q) = g_{\mu\nu} - q_\mu q_\nu / q^2 = g_{\mu\nu} - L_{\mu\nu}(q)$. Its free counterpart $D_0 \equiv D_{\mu\nu}^0(q)$ is obtained by putting the full gluon form factor $d(-q^2, \xi)$ in Eq. (2.2) simply to one, i.e., $D_{\mu\nu}^0(q) = - \{ T_{\mu\nu}(q) + \xi L_{\mu\nu}(q) \} (1/q^2)$. In order to evaluate the effective potential (2.1), on account of Eq. (2.2), we use the well-known expression, namely

$$\text{Tr} \ln(D_0^{-1} D) = 8 \times 4 \ln \det(D_0^{-1} D) = 32 \ln[(3/4) d(-q^2, \xi) + (1/4)]. \quad (2.3)$$

It becomes zero (in accordance with the above-mentioned normalization condition) when the full gluon form factor is replaced by its free counterpart.

Going over to four-dimensional Euclidean space in Eq. (2.1), and evaluating some numerical factors, one obtains ($\epsilon_g = V(D)$)

$$\epsilon_g = - \frac{1}{\pi^2} \int dq^2 q^2 \left[\ln[1 + 3d(q^2, \xi)] - \frac{3}{4} d(q^2, \xi) + a \right], \quad (2.4)$$

where constant $a = (3/4) - 2 \ln 2 = -0.6363$ and the integration from zero to infinity is assumed. The VED ϵ_g derived in Eq. (2.4) is already colorless quantity, since it has been already summed over color indices. It does not also depend explicitly on the unphysical (longitudinal) part of the full gluon propagator due to the product $(D_0^{-1} D)$, which in its turn comes from the above-mentioned normalization to zero (see above).

However, it still suffers from the two serious problems. The coefficient of the transversal Lorentz structure (the effective charge $d(q^2, \xi)$) may still depend explicitly on ξ . Also, it is badly divergent at least as the fourth power of the ultraviolet (UV) cutoff, and therefore suffers from different types of the PT contributions (see below).

III. THE TRULY NP VED

In order to define the VED free of all the above-mentioned problems, let us make first the identical transformation of the full gluon form factor (which is nothing but the effective charge) in Eq. (2.2) as follows (Euclidean signature already):

$$d(q^2, \xi) = d(q^2, \xi) - d^{PT}(q^2, \xi) + d^{PT}(q^2, \xi) = d^{NP}(q^2, \xi) + d^{PT}(q^2, \xi), \quad (3.1)$$

where $d^{PT}(q^2, \xi)$ correctly describes the PT structure of the effective charge $d(q^2, \xi)$, including its behavior in the UV limit (asymptotic freedom [12]), otherwise remaining arbitrary. On the other hand, $d^{NP}(q^2, \xi)$, defined by the

above-made subtraction, is assumed to correctly reproduce the NP structure of the effective charge, including its asymptotic in the deep infrared (IR) limit, underlying thus the strong intrinsic influence of the IR properties of the theory on its NP dynamics. Evidently, both terms are valid in the whole energy/momentum range, i.e, they are not asymptotics. Let us also emphasize the principle difference between $d(q^2, \xi)$ and $d^{NP}(q^2, \xi)$. The former is the NP quantity "contaminated" by the PT contributions, while the latter one being also NP, nevertheless, is free of them. Thus the exact separation between the truly NP effective charge $d^{NP}(q^2, \xi)$ and its PT counterpart $d^{PT}(q^2, \xi)$ is achieved (for the concrete example considered below this separation is unique as well).

There is also another serious reason for the subtraction (3.1). The problem is that the UV asymptotic of the full effective charge due to asymptotic freedom may depend on the gauge-fixing parameter, namely to leading order

$$d(q^2, \xi) \sim_{q^2 \rightarrow \infty} (\ln(q^2/\Lambda_{QCD}^2))^{c_0/b_0}, \quad (3.2)$$

where the exponent c_0/b_0 explicitly depends on the gauge-fixing parameter via the coefficient c_0 [12], and Λ_{QCD}^2 is the QCD asymptotic scale parameter. Evidently, in the decomposition (3.1) precisely the PT part of the full effective charge will be responsible for this explicit dependence on the gauge choice. Subtracting it, we will be guaranteed that the remaining part will not explicitly depend on the gauge-fixing parameter (see below).

Substituting the exact decomposition (3.1) into Eq. (2.4) and doing some trivial rearrangement, one obtains

$$\epsilon_g = -\frac{1}{\pi^2} \int dq^2 q^2 \left[\ln[1 + 3d^{NP}(q^2, \xi)] - \frac{3}{4}d^{NP}(q^2, \xi) \right] + \epsilon_{PT}, \quad (3.3)$$

where we introduce the following notation

$$\epsilon_{PT} = -\frac{1}{\pi^2} \int dq^2 q^2 \left[\ln\left[1 + \frac{3d^{PT}(q^2, \xi)}{1 + 3d^{NP}(q^2, \xi)}\right] - \frac{3}{4}d^{PT}(q^2, \xi) + a \right]. \quad (3.4)$$

It contains the contribution which is mainly determined by the PT part of the full gluon propagator, $d^{PT}(q^2, \xi)$. The constant a also should be included, since it comes from the normalization of the free PT vacuum to zero. However, this is not the whole story yet. The first term in Eq. (3.3), depending only on the truly NP effective charge, nevertheless, assumes the integration over the PT region (up to infinity), which should be subtracted as well. The problem is that the first term in Eq. (3.3) may still be divergent in the UV limit. *A priori*, there is only one restriction on its UV asymptotic, namely do not contradict to asymptotically free behavior of the full effective charge, to leading order shown in Eq. (3.2). For example, if $d^{NP}(q^2, \xi)$ is a linearly falling at infinity function, then the integral in Eq. (3.3) still diverges as mentioned above.

If we separate the NP region from the PT one, by introducing the so-called effective scale q_{eff}^2 explicitly, then we get

$$\epsilon_g = -\frac{1}{\pi^2} \int_0^{q_{eff}^2} dq^2 q^2 \left[\ln[1 + 3d^{NP}(q^2, \xi)] - \frac{3}{4}d^{NP}(q^2, \xi) \right] + \epsilon_{PT} + \epsilon'_{PT}, \quad (3.5)$$

where evidently

$$\epsilon'_{PT} = -\frac{1}{\pi^2} \int_{q_{eff}^2}^{\infty} dq^2 q^2 \left[\ln[1 + 3d^{NP}(q^2, \xi)] - \frac{3}{4}d^{NP}(q^2, \xi) \right]. \quad (3.6)$$

This integral represents contribution to the VED which is determined by the truly NP part of the full gluon propagator but integrated out over the PT region. Along with ϵ_{PT} given in Eq. (3.4) it also represents a type of the PT contribution into the gluon part of the VED (3.5). This means that the two remaining terms in Eq. (3.5) should be subtracted by introducing the truly NP YM VED ϵ_{YM} as follows:

$$\epsilon_{YM} = \epsilon_g - \epsilon_{PT} - \epsilon'_{PT}, \quad (3.7)$$

where the explicit expression for ϵ_{YM} is given by the integral in Eq. (3.5) (see below as well). Concluding this section, let us only note that the above-mentioned necessary subtractions can be done in a more sophisticated way by means of ghost degrees of freedom [9]. In other words, the ghost degrees of freedom are to be included into the PT parts of the VED. Evidently, the subtracted terms are of no importance for our present consideration.

IV. THE BAG CONSTANT

The Bag constant (the so-called the Bag pressure) is defined as the difference between the PT and the NP VED [13]. So in our notations for YM fields, and as it follows from the definition (3.7), it is nothing but the truly NP YM VED apart from the sign, i.e.,

$$\begin{aligned} B_{YM} = -\epsilon_{YM} &= \epsilon_{PT} + \epsilon'_{PT} - \epsilon_g \\ &= \frac{1}{\pi^2} \int_0^{q_{eff}^2} dq^2 q^2 \left[\ln[1 + 3d^{NP}(q^2, \xi)] - \frac{3}{4}d^{NP}(q^2, \xi) \right]. \end{aligned} \quad (4.1)$$

This is a general expression for any model effective charge in order to calculate the Bag constant (or the truly NP YM VED apart from the sign) from first principles. It is our definition of the truly NP YM VED and thus of the Bag constant (apart from the sign) as integrated out the truly NP effective charge over the NP region (soft momentum region, $0 \leq q^2 \leq q_{eff}^2$). It is free of all types of the PT contributions, by construction. In this connection, let us recall that ϵ_g is also NP, but "contaminated" by the PT contributions, which just to be subtracted in order to get expression (4.1).

Comparing expressions (2.4) and (4.1), one comes to the following *prescription* to derive Eq. (4.1) directly from Eq. (2.4).

- (i). Replacing $d(q^2) \rightarrow d^{NP}(q^2)$.
- (ii). Omitting the constant a which normalizes the free PT vacuum to zero.
- (iii). Introducing the effective scale q_{eff}^2 which separates the NP region from the PT one.
- (iv). Omitting the minus sign for the Bag constant.

At this stage the Bag constant (4.1) is free of all types of the PT "contaminations", and it is finite quantity, for sure. All other its properties mentioned above (positivity, no imaginary part, etc.) depend on the chosen effective charge. It is worth emphasizing once more that the Bag constant and hence the truly NP VED is the main physical characteristic of the true QCD ground state. Let us note that in defining correctly the truly NP YM VED (or equivalently the Bag constant), the three types of the corresponding subtractions have been introduced. The first one - in Eq. (3.1) at the fundamental gluon level and the two others - in Eq. (3.7). For the general discussion of necessity in such types of the subtractions see Ref. [14].

The quantum part of the total truly NP VED at log-loop level is

$$\epsilon_t = \epsilon_{YM} + N_f \epsilon_q, \quad (4.2)$$

where ϵ_q is the truly NP quark loop contribution. It is an order of magnitude less than ϵ_{YM} because of much less quark degrees of freedom in the vacuum, and it is positive because of overall minus due to quark loop. Evidently, in terms of the Bag constant, one obtains

$$\epsilon_t = -B_{YM}[1 - N_f \nu], \quad (4.3)$$

where we introduce $\epsilon_q = \nu B_{YM}$ and $\nu \ll 1$.

V. NUMERICAL EVALUATION OF THE BAG CONSTANT

Eq. (4.1) is the main subject of our consideration in what follows. The only problem remaining to solve is to choose such truly NP effective charge $d^{NP}(q^2, \xi)$ which should not *explicitly* depend on the gauge-fixing parameter ξ . Here it is worth emphasizing that the implicit gauge dependence is not problem. Such kind of the dependence is unavoidable in gauge theories like QCD, since fields themselves are gauge-dependent [15]. For different truly NP effective charges $d^{NP}(q^2, \xi)$ we will get different numerical results. That is why the choice for its explicit expression should be physically and mathematically well justified.

For concrete numerical evaluation of the Bag constant (4.1), let us choose the truly NP effective charge as follows:

$$d^{NP}(q^2) \equiv \alpha_s^{NP}(q^2) = \Lambda_{NP}^2/q^2, \quad (5.1)$$

where Λ_{NP} is the mass scale parameter (the mass gap) responsible for the large-scale structure of the true QCD vacuum. First of all, in continuous QCD it is the NP solution of the Schwinger-Dyson (SD) equation for the full gluon

propagator [14] (and references therein). It is well known that in this theory it leads to the linear rising potential between heavy quarks, "seen" by lattice QCD [16] as well $((q^2)^{-2}$ -type behavior for the full gluon propagator). Secondly, it does not depend explicitly on the gauge choice. Also, the separation between the truly NP and PT effective charge is not only exact but is unique as mentioned above, since the PT effective charge is always regular at zero, while the truly NP effective charge (5.1) is singular at origin. Let us note in advance that it leads to many other desirable properties for the Bag constant (see below).

In order to proceed to the explicit numerical calculation of the Bag constant for the truly NP effective charge (5.1), let us introduce now dimensionless variable and parameter as follows:

$$z = q^2/\Lambda_{NP}^2, \quad z_c = q_{eff}^2/\Lambda_{NP}^2, \quad d^{NP}(q^2) = d^{NP}(z) = 1/z. \quad (5.2)$$

From the general expression for the Bag constant (4.1) one then gets the dimensionless truly NP YM effective potential defined at a fixed effective scale squared q_{eff}^4 as follows:

$$\Omega_{YM}(z_c) = \frac{1}{q_{eff}^4} \epsilon_{YM}(q_{eff}^4, z_c) = \frac{1}{\pi^2} z_c^{-2} \int_0^{z_c} dz \, z \left[\frac{3}{4} d^{NP}(z) - \ln[1 + 3d^{NP}(z)] \right], \quad (5.3)$$

which obviously makes it possible to factorize the scale dependence in the truly NP YM VED (4.1). Performing almost trivial integration in this expression, one obtains

$$\Omega_{YM}(z_c) = \frac{1}{2\pi^2} z_c^{-2} \left[9 \ln \left(1 + \frac{z_c}{3} \right) - \frac{3}{2} z_c - z_c^2 \ln \left(1 + \frac{3}{z_c} \right) \right]. \quad (5.4)$$

It is easy to see now that as a function of z_c , the effective potential (5.4) diverges as $\sim z_c^{-1}$ at small z_c (the unphysical regime, $\Lambda_{NP} \rightarrow \infty$ because the mass gap is either finite or zero, i.e., it cannot be infinitely large). It converges as $\sim -z_c^{-1}$ at infinity (PT limit, $\Lambda_{NP} \rightarrow 0$), i.e. it approaches zero from below. Let us emphasize that in the PT limit the truly NP phase vanishes and the PT phase survives only. At a fixed effective scale q_{eff}^2 it follows that $z_c \rightarrow \infty$, indeed. In other words, at a fixed effective scale one recovers correct PT limit, while at a fixed mass gap it evidently cannot be recovered.

From the above one can conclude that the effective potential (5.4) as a function of z_c has a minimum at some finite point (local minimum), so its minimization makes sense. In general, by taking first derivatives of the effective potential one recovers the corresponding equations of motion [10, 11]. Thus the above-mentioned minimization of the effective potential (5.4) makes it possible to fix the constant of integration which is z_c in our case. Requiring $\partial\Omega_{YM}(z_c)/\partial z_c = 0$, yields the following "stationary" condition

$$z_c = 4 \ln[1 + (z_c/3)]. \quad (5.5)$$

Its numerical solution is

$$z_c^{min} = 2.2, \quad (5.6)$$

so at the "stationary" state the effective potential (5.3) can be written down as follows:

$$\Omega_{YM}(z_c^{min}) = \frac{1}{2\pi^2} \left[\frac{3}{4} (z_c^{min})^{-1} - \ln \left(1 + \frac{3}{z_c^{min}} \right) \right] = -0.0263. \quad (5.7)$$

The truly NP YM VED (5.3) and hence the Bag constant thus becomes

$$B_{YM} = -\epsilon_{YM} = 0.0263 q_{eff}^4 = 0.1273 \times \Lambda_{NP}^4, \quad (5.8)$$

where the relation

$$q_{eff}^2 = z_c^{min} \Lambda_{NP}^2 = 2.2 \Lambda_{NP}^2 \quad (5.9)$$

has been already used.

So, we have explicitly demonstrated that the Bag constant (5.8) at the "stationary" state is finite, positive, and it has no imaginary part, indeed. It depends only on the mass gap responsible for the truly NP dynamics in the QCD ground state, or equivalently on the effective scale separating the NP region from the PT one. Evidently, only the minimization of the effective potential (5.4) makes it possible to fix exactly the numerical relation (5.9) between them.

VI. NUMERICAL RESULTS

In order to complete the numerical calculation of the Bag constant (5.8) all we need now is the concrete value for the effective scale q_{eff} , which separates the NP region from the PT one. Equivalently, the concrete value for a scale at which the NP effects become important, that is the mass gap Λ_{NP} , also allows one to achieve the same goal.

If the PT region starts conventionally from 1 GeV , then this number is a natural choice for the effective scale. It makes it also possible to directly compare our values with the values of many phenomenological parameters calculated just at this scale (see Appendix A below). We consider this value as well justified and realistic upper limit for the effective scale defined above. Thus, we put

$$q_{eff} = q_{eff}^2 = 1 \text{ GeV}, \quad (6.1)$$

then for the mass gap, on account of the relation (5.8), we get

$$\Lambda_{NP} = 0.6756 \text{ GeV}. \quad (6.2)$$

Equivalently, the numerical value of the mass gap Λ_{NP} has been obtained from the experimental value for the pion decay constant, $F_\pi = 93.3 \text{ MeV}$, by implementing a physically well-motivated scale-setting scheme [17, 18]. The pion decay constant is good experimental number, since it is directly measured quantity in comparison, for example with the quark or the gluon condensates, etc. For the mass gap we have obtained the following numerical result:

$$\Lambda_{NP} = 0.5784 \text{ GeV}, \quad (6.3)$$

which for the effective scale yields

$$q_{eff} = 0.857 \text{ GeV}, \quad q_{eff}^2 = 0.735 \text{ GeV}^2. \quad (6.4)$$

In what follows we will consider this value as a realistic lower limit for the effective scale. One has to conclude that we have obtained rather close numerical results for the effective scale and the mass gap, by implementing rather different scale-setting schemes. It is worth emphasizing that the effective scale (6.4) quite well covers not only the deep IR region but the substantial part of the intermediate one as well.

For the Bag constant (and hence for the truly NP YM VED) from Eq. (5.8), one obtains

$$\begin{aligned} B_{YM} = -\epsilon_{YM} &= (0.0142 - 0.0263) \text{ GeV}^4 \\ &= (1.84 - 3.4) \text{ GeV}/fm^3 = (1.84 - 3.4) \times 10^{39} \text{ GeV}/cm^3. \end{aligned} \quad (6.5)$$

This is a huge amount of energy stored in one cm^3 of the NP QCD vacuum even in "God-given" units $\hbar = c = 1$. In order to restore the explicit dependence on $\hbar$ [10], the right-hand-side of Eq. (6.5) should be multiplied by it, which numerical value in different units is $\hbar = 1 \times 10^{-34} \text{ J s} = 6.6 \times 10^{-25} \text{ GeV s}$. Evidently, using the number of different conversion factors (see, for example Ref. [15] or Particle Data Group) the Bag constant can be expressed in different systems of units (SI, CGS, etc.).

Taking into account that

$$1 \text{ GeV} = 0.15 \times 10^{-9} \text{ J} = 0.48 \times 10^{-17} \text{ W}, \quad (6.6)$$

from Eq. (6.5) one finally gets ($1 \text{ W} = 10^{-3} \text{ kW} = 10^{-6} \text{ MW} = 10^{-9} \text{ GW}$)

$$B_{YM} = (0.88 - 1.63) \times 10^{13} \text{ GW}/cm^3, \quad (6.7)$$

in familiar units of watt (W). This number still indicates a huge amount of energy stored in one cm^3 of the NP QCD vacuum especially in comparison with the total nuclear energy produced in one year and which is estimated about 360 GW only.

VII. DISCUSSION

We have already pointed out that the VED is badly divergent as the fourth power of the UV cutoff, so we can write

$$\epsilon_g = -\Lambda_{UV}^4 = -\lambda^4 B, \quad (7.1)$$

which can be always done without losing generality, and where λ is the dimensionless UV cutoff. At the same time, we have obtained that the truly NP VED is finite and it is $\epsilon_{YM} = -B$. This means that the further analysis can be done in terms of the Bag constant.

Let us imagine now that we can extract the finite portion ϵ_{YM} from the vacuum in k different places (different "vacuum energy releasing facilities" (VERF)). It can be done by n_m times in each place, where $m = 1, 2, 3 \dots k$. Then the remaining (R) VED becomes

$$\epsilon_R = \epsilon_g - \epsilon_E = \epsilon_g - \sum_{m=1}^k n_m \epsilon_{YM} = \epsilon_g + B \sum_{m=1}^k n_m, \quad (7.2)$$

where, evidently, ϵ_E denotes the extracting (E) VED.

The ideal case (which, however, will never be achieved) is when we could extract the finite portion of the VED an infinitely number times and in an infinitely number places, so we should consider the following limits:

$$\epsilon_R = B \lim_{(\lambda, k, n_m) \rightarrow \infty} \left[-\lambda^4 + \sum_{m=1}^k n_m \right]. \quad (7.3)$$

Thus, one can conclude that in the ideal case

$$\epsilon_E = B \lim_{(\lambda, k, n_m) \rightarrow \infty} \sum_{m=1}^k n_m \sim B \times \lambda^2, \quad (7.4)$$

diverges quadratically since due to the limits pointed out in Eq. (7.3), we can approximate $k \sim \lambda$, $n_m \sim \lambda$, $\lambda \rightarrow \infty$. At the same time, the initial VED ϵ_g diverges quartically, i.e., $\epsilon_g \sim -B \times \lambda^4$, $\lambda \rightarrow \infty$ (see Eq. (7.1)). This means that the remaining VED

$$\epsilon_R = \epsilon_g - \epsilon_E \sim B \lim_{\lambda \rightarrow \infty} [-\lambda^4 + \lambda^2], \quad (7.5)$$

will be always equal to the initial VED ϵ_g in the limit pointed out above, i.e., in units of B one has

$$\epsilon_R = \epsilon_g - \epsilon_E = \epsilon_g + O(1/\lambda^2), \quad \lambda \rightarrow \infty. \quad (7.6)$$

This will be especially so in the real case when the integer numbers k and n_m are very big, but finite. The true QCD vacuum is an infinite and permanent source of energy. The only problem is how to extract it in the profitable way. This is, of course, beyond the scope of this paper, though there are some preliminary ideas on this and related issues.

It is instructive to show explicitly that the same relation (7.6) takes place between not only the different VED's but between different types of energy themselves. Indeed, the different VED's should be multiplied by volume (V) in order to get energy. Then from Eqs. (6.5) and (7.4) in units of $[(1.84 - 3.4) \text{ GeV}]$ it follows that

$$E_E = V \epsilon_E = -(1.84 - 3.4) \text{ GeV} \frac{V}{fm^3} \times \lim_{(\lambda, k, n_m) \rightarrow \infty} \sum_{m=1}^k n_m \sim -\lambda^5, \quad (7.7)$$

since $V/fm^3 \sim \lambda^3$ always when λ goes to infinity.

In the same units, one gets

$$E_g = V\epsilon_g = -(1.84 - 3.4) \text{ GeV} \frac{V}{fm^3} \times \lambda^4 \sim -\lambda^7. \quad (7.8)$$

So the difference between them which is nothing but the remaining energy E_R becomes

$$E_R = E_g - E_E = E_g + O(1/\lambda^2), \quad \lambda \rightarrow \infty, \quad (7.9)$$

in complete agreement with the relation (7.6) as it should be. Let us remind that $[(1.84 - 3.4) \text{ GeV}]$ units are equal to $[B \text{ fm}^3]$ units, so in fact this simple analysis is again performed in terms of the Bag constant.

VIII. CONCLUSIONS

In summary, we have formulated a general method as to how to numerically calculate the quantum part of the truly NP YM VED (the Bag constant, apart from the sign, by definition) in the covariant gauge QCD ground state, using the effective potential approach for composite operators. The Bag constant is defined as integrated out the truly NP part of the effective charge over the NP region (soft momentum region), Eq. (4.1). At this general stage the Bag constant is colorless (color-singlet), finite, and it is free of all types of the PT contributions, by construction. The separation of "truly NP versus PT" contributions into the VED is exact and unique because of the subtraction at the fundamental level (3.1), as well as due to all other subtractions explicitly shown in Eq. (3.7).

For the considered truly NP effective charge (5.1) in addition it is negative, and it has no imaginary part (stable vacuum). It is also a manifestly gauge-invariant quantity (i.e., does not explicitly depend on the gauge-fixing parameter as it is required). The separation of "soft versus hard" gluon momenta is also exact because of the minimization procedure. It becomes possible since the effective potential (5.4) as a function of the dimensionless effective scale z_c , which separates the NP region from the PT one, has a local minimum. Our numerical value for the Bag constant is in a good agreement with other NP quantities such as the gluon condensate. In order to demonstrate this we have used the trace anomaly relation, not applying to the weak coupling solution for the β function (see Appendix A).

Our method can be generalized on the multi-loop contributions in the effective potential, as well as to take into account the quark degrees of freedom. These terms, however, will produce numerically the contributions an order of magnitude less, at least, in comparison with the leading log-loop level gluon term. What is necessary indeed, is to be able to extract the finite part of the truly NP VED in a self-consistent and manifestly gauge-invariant ways. Just this is provided by our method which thus can be applied to any QCD vacuum quantum and classical models. It may serve as a test of them, providing an exact criterion for the separation "stable versus unstable" vacua. Using our method we have already shown that the vacuum of classical dual Abelian Higgs model with string and without string contributions is unstable against quantum corrections [19, 20].

The Bag constant calculated here is a manifestly gauge-invariant and colorless finite quantity, i.e., it is a physical quantity. That is why it makes sense to discuss its releasing from the vacuum. That's the VED in general is badly divergent is not a mathematical problem. This reflects an universal reality. Vacuum is everywhere and it always exists. Our Universe in general and our real world in particular is only its special type of excitation due to the Big Bang. As underlined above, the vacuum is infinite and hence permanent source of energy. The only problem is how to release the finite portion (the Bag constant) and whether it will be profitable or not by introducing some type of cyclic process. "Perpetuum mobile" does not exist, but "perpetuum source" of energy does exist and it is the QCD ground state. Analogously to the Plank constant $\hbar$ which determines a quant of energy emitted by quantum physical systems, the Bag constant determines a quant of energy which can be released from the QCD vacuum.

A financial support from HAS-JINR Scientific Collaboration Fund and Hungarian OTKA-T043455 (P. Levai) is to be acknowledged.

APPENDIX A: THE TRACE ANOMALY RELATION

The truly NP VED (and hence the Bag constant) is important in its own right as the main characteristic of the QCD ground state. Furthermore it assists in calculating such an important phenomenological parameter as the gluon condensate, introduced in the QCD sum rules approach to resonance physics [21]. The famous trace anomaly relation [22] in the general case (nonzero current quark masses m_f^0) is

$$\Theta_{\mu\mu} = \frac{\beta(\alpha_s)}{4\alpha_s} G_{\mu\nu}^a G_{\mu\nu}^a + \sum_f m_f^0 \bar{q}_f q_f. \quad (A1)$$

where $\Theta_{\mu\mu}$ is the trace of the energy-momentum tensor and $G_{\mu\nu}^a$ being the gluon field strength tensor while $\alpha_s = g^2/4\pi$. Sandwiching Eq. (A.1) between vacuum states and on account of the obvious relation $\langle 0|\Theta_{\mu\mu}|0\rangle = 4\epsilon_t$, one obtains

$$4\epsilon_t = \langle 0|\frac{\beta(\alpha_s)}{4\alpha_s}G_{\mu\nu}^a G_{\mu\nu}^a|0\rangle + \sum_f m_f^0 \langle 0|\bar{q}_f q_f|0\rangle. \quad (\text{A2})$$

Here ϵ_t is the sum of all possible independent, NP contributions to the VED (the total VED) and $\langle 0|\bar{q}_f q_f|0\rangle$ is the chiral quark condensate. From this equation in the case of the pure YM fields (i.e., when the number of quark fields is zero $N_f = 0$), one obtains

$$\langle 0|\frac{\beta(\alpha_s)}{4\alpha_s}G_{\mu\nu}^a G_{\mu\nu}^a|0\rangle = 4\epsilon_{YM}, \quad (\text{A3})$$

where, evidently, we saturate the total VED ϵ_t by the truly NP YM VED ϵ_{YM} defined in Eq. (4.1) and calculated in Eq. (6.5) at log-loop level, i.e., setting $\epsilon_t = \epsilon_{YM} + \dots$. If confinement happens then the β function is always in the domain of attraction (i.e., always negative) without IR stable fixed point [12]. Therefore it is convenient to introduce the general definition for the gluon condensate (i.e., without using the weak coupling limit solution to the β function) as follows:

$$\langle G^2 \rangle \equiv -\langle 0|\frac{\beta(\alpha_s)}{4\alpha_s}G_{\mu\nu}^a G_{\mu\nu}^a|0\rangle = -4\epsilon_{YM} = 4B_{YM}. \quad (\text{A4})$$

Such defined general gluon condensate will be always positive as it should be.

The renormalization group equation for the β function

$$q^2 \frac{d\alpha_s(q^2)}{dq^2} = \beta(\alpha_s(q^2)) \quad (\text{A5})$$

after substitution of our solution for the truly NP effective charge (5.1) yields

$$\beta(\alpha_s(q^2)) = -\alpha_s(q^2), \quad (\text{A6})$$

so that $\beta(\alpha_s)/\alpha_s = \beta(\alpha_s(q^2))/\alpha_s(q^2) = -1$, and Eq. (A.4) numerically becomes

$$\langle G^2 \rangle \equiv -\langle 0|\frac{\beta(\alpha_s)}{4\alpha_s}G_{\mu\nu}^a G_{\mu\nu}^a|0\rangle = \langle 0|\frac{1}{4}G_{\mu\nu}^a G_{\mu\nu}^a|0\rangle = 4B_{YM} = (0.0568 - 0.1032) \text{ GeV}^4, \quad (\text{A7})$$

where we have used the numerical value for the Bag constant (6.5). To the gluon condensate (A.7) it can be assigned a physical meaning indeed as the global (average) vacuum characteristic which measures a density of the NP gluon fields configurations in the true QCD vacuum.

However, it cannot be directly compared with the phenomenological values for the standard gluon condensate estimated within different approaches [23]. The problem is that it is necessary to remember that any value at the scale Λ_{NP} (lower bound in the right-hand-side of Eq. (A.7)) is to be recalculated at the 1 GeV scale. Moreover, both values explicitly shown in Eq. (A.7) should be recalculated at the same parametrization, which is nothing but the ratio $\beta(\alpha_s)/\alpha_s$. In phenomenology the standard parametrization for the gluon condensate is

$$G_2 = \langle \frac{\alpha_s}{\pi} G^2 \rangle = \langle 0|\frac{\alpha_s}{\pi} G_{\mu\nu}^a G_{\mu\nu}^a|0\rangle \approx 0.012 \text{ GeV}^4, \quad (\text{A8})$$

which can be changed within a factor of 2 [19].

Thus in order to achieve the same parametrization the both sides of Eq. (A.7) should be multiplied by $4(\alpha_s/\pi)$. For the numerical value of the strong fine structure constant we use $\alpha_s = \alpha_s(m_Z) = 0.1187$ from the Particle Data Group. In addition the lower bound should be multiplied by the factor $1 \text{ GeV}/\Lambda_{NP} = 1.57$, coming from the numerical value (6.3). Then the recalculated gluon condensate (A.7), which is denoted as $\bar{G}_2$, finally becomes

$$\bar{G}_2 = (0.014 - 0.022) \text{ GeV}^4. \quad (\text{A9})$$

Just this numerical value for the gluon condensate should be compared with the numerical value coming from the phenomenology, see Eq. (A.8) above. So there is no doubt that all our numerical results are in good agreement with other phenomenological estimates [23].

-
- [1] K. A. Milton, hep-th/0406024.
 - [2] M. D. Roberts, hep-th/0012063;
G. L. Klimchitskaya, V. M. Mostepanenko, quant-ph/0609145.
 - [3] S.-S. Xue, Phys. Lett. B 508 (2001) 211.
 - [4] S.-S. Xue, Phys. Rev. D 68 (2003) 013004.
 - [5] M. Scandurra, hep-th/0104127.
 - [6] Confinement, Duality, and Nonperturbative Aspects of QCD, edited by P. van Baal, NATO ASI, Series B: Physics, vol. 368 (Plenum, New York, 1997).
 - [7] Non-Perturbative QCD, Structure of the QCD Vacuum, edited by K-I. Aoki, O. Miymura, and T. Suzuki [Prog. Theor. Phys. Suppl. 131 (1998) 1].
 - [8] T. Schafer, E. V. Shuryak, Rev. Mod. Phys. 70 (1998) 323;
V. A. Rubakov, Classical Thory of Gauge Fields (Princeton University Press, 2002).
 - [9] V. Gogohia, Gy. Kluge, Phys. Rev. D 62 (2000) 076008.
 - [10] J. M. Cornwall, R. Jackiw, E. Tomboulis, Phys. Rev. D 10 (1974) 2428.
 - [11] A. Barducci et al., Phys. Rev. D 38 (1988) 238;
R. W. Haymaker, Riv. Nuovo Cim. 14 (1991) 1-89.
 - [12] W. Marciano, H. Pagels, Phys. Rep. C 36 (1978) 137.
 - [13] E. V. Shuryak, Phys. Rep. 115 (1984) 151;
M. S. Chanowitz, S. Sharpe, Nucl. Phys. B 222 (1983) 211.
 - [14] V. Gogokhia, hep-ph/0606060.
 - [15] M. E. Peskin, D. V. Schroeder, An Introduction to Quantum Field Theory (AW, Advanced Book Program, 1995).
 - [16] K. D. Born et al., Phys. Lett. B 329 (1994) 325;
V. M. Miller et al., Phys. Lett. B 335 (1994) 71.
 - [17] V. Gogohia, Gy. Kluge, H. Toki, T. Sakai, Phys. Lett. B 453 (1999) 281.
 - [18] V. Gogohia, H. Toki, T. Sakai, Gy. Kluge, Inter. Jour. Mod. Phys. A 15 (2000) 45.
 - [19] V. Gogohia, Gy. Kluge, Phys. Lett. B 477 (2000) 387.
 - [20] V. Gogohia, M. Prisznyak, Phys. Lett. B 494 (2000) 109.
 - [21] M. A. Shifman, A. I. Vainshtein, V. I. Zakharov, Nucl. Phys. B 147 (1979) 385.
 - [22] R. J. Crewther, Phys. Rev. Lett. 28 (1972) 1421;
M. S. Chanowitz, J. Ellis, Phys. Rev. D 7 (1973) 2490;
J. C. Collins, A. Duncan, S. D. Joglecar, Phys. Rev. D 16 (1977) 438.
 - [23] S. Narison, Phys. Lett. B 387 (1996) 162.